



# Antioxidants Against Free Radicals

©2008 Huntington College of Health Sciences

Literature Education Series On Dietary Supplements

By Gene Bruno, MS, MHS

*Smart Supplementation™ is a free series of educational literature created by Huntington College of Health Sciences (HCHS) as a public service. Although copyrighted, it may be freely photocopied and distributed, but may not be altered in any way. Smart Supplementation™ is not intended as medical advice. For diagnosis and treatment of any medical condition, consult your physician.*

Free radicals are atoms with one or more unpaired electrons, which creates an imbalance in the way the atoms function. They are formed when oxygen interacts with certain molecules. Once formed these highly reactive radicals can start a chain reaction which results in damage to important cellular components such as DNA, or the cell membrane. Cells may function poorly or die if this occurs. To prevent free radical damage the body has a defense system of antioxidants.

Antioxidants protect against cellular damage caused by free radicals, which would otherwise be the common pathway for cancer, aging, and a variety of diseases. Following is a discussion about some of the most common and valuable antioxidants that we can use in supplemental form to protect ourselves.

## VITAMIN A

Beta-carotene is perhaps best known for the fact that the body can convert it into vitamin A. This takes place in the intestinal wall (although some diabetics have problems making this conversion). Since beta carotene is only converted to vitamin A as the body needs it, it is virtually impossible for beta carotene to cause vitamin A toxicity.

### Antioxidant effects

In addition to functioning as a vitamin A precursor, beta-carotene has also been shown to provide other significant health benefits. Beta-carotene has antioxidant activities and prevents lipid peroxidation.<sup>1</sup> Serum beta-carotene levels are inversely related to C-reactive protein levels and the white blood count. These markers are markers of

inflammation and are associated with suboptimal cardiovascular function.<sup>2</sup> The antioxidant effects of beta-carotene may reduce free radical-induced DNA damage.<sup>3, 4</sup>

### Reduced risk of mutated lung cell growth

In addition, as intake of beta-carotene from food increases, the risk of mutated lung cell growth is reduced<sup>5</sup>. It is possible that similar benefits may be realized from supplementation of natural beta-carotene. Beta-carotene has also been shown to be effective in reducing the risk of pre-mutated cells the mouth. In one study<sup>6</sup>, people with such cells received 180 mg of beta-carotene twice weekly. After 6 months, the results were a reduced frequency of the mutated cells, and inhibition of new lung cell mutations. In a similar study<sup>7</sup>, 60 mg twice daily of beta-carotene was also found to be effective for the same purpose.

### Does beta-carotene increase the risk of lung cancer in smokers?

No, it does not. This misconception was based upon one flawed study in which smokers using beta-carotene were thought to be at a greater risk of lung cancer (*N Engl J Med* 1994; 330:1029-35). However, a more thorough follow-up analysis which looked at the diets and other dietary supplements taken, revealed that the smokers' actual danger was due to low total antioxidant levels, not to the fact that they took beta-carotene (*Am J Epidemiol* 2004; 160(1):68-76). This makes sense given the fact that antioxidants function interdependently, and so should be taken together.

## VITAMIN C

Vitamin C is probably best known for its effects as an antioxidant and its role in maintaining proper immune function.<sup>8</sup> Many of the beneficial effects of vitamin C are attributed primarily to antioxidant and free radical scavenging effects. Vitamin C readily undergoes reversible oxidation and reduction in the body.<sup>9</sup> Vitamin C decreases oxidants in gastric juice, lipid peroxidation, and oxidative DNA and protein damage.<sup>10</sup> Vitamin C might reduce oxidative stress caused by the large number of oxidants in cigarette smoke.<sup>11</sup>

## VITAMIN E

The term vitamin E describes a family of eight antioxidants: four tocopherols and four tocotrienols. Alpha-tocopherol is the only form of vitamin E that is actively maintained in the human body; therefore, it is the form of vitamin E found in the largest quantities in blood and tissues.<sup>12</sup> It is also the only form that meets the latest Recommended Dietary Allowance (RDA) for vitamin E. Nevertheless the other tocopherols and tocotrienols have significant contributions to make toward human health. These other tocopherols include the beta, gamma, and delta tocopherols; and are considered isomers. In this context, the term isomer refers to compounds with the same structural formula but different spatial arrangements of atoms.<sup>13</sup> In addition, there is a family of natural compounds related to vitamin E called tocotrienols, which also have distinct health benefits. These tocotrienols are also found in alpha, beta, delta and gamma isomers.

### Alpha-tocopherol

The main function of alpha-tocopherol in humans appears to be that of an antioxidant. Free radicals are formed primarily in the body during normal metabolism and also upon exposure to environmental factors, such as cigarette smoke or pollutants. Fats, which are an integral part of all cell membranes, are vulnerable to destruction through oxidation by free radicals. The fat-soluble vitamin, alpha-tocopherol, is uniquely suited to intercept free radicals and thus prevent a chain reaction of lipid destruction. Aside from maintaining the integrity of cell membranes throughout the body, alpha-tocopherol also protects the fats in low density lipoproteins (LDLs) from oxidation. Lipoproteins are particles composed of lipids and proteins that transport fats through the bloodstream. LDLs specifically transport cholesterol from the liver to the tissues of the body. Oxidized LDLs have been implicated in the development of cardiovascular diseases. When a molecule of alpha-tocopherol neutralizes a free radical, it is altered in such a way that its antioxidant capacity is lost. However, other antioxidants, such as vitamin C, are capable of regenerating the antioxidant capacity of alpha-tocopherol.<sup>14 15</sup>

Several other functions of alpha-tocopherol have been identified that are not likely related to its antioxidant capacity. For instance, alpha-tocopherol appears to also affect the expression and activities of molecules and enzymes in immune and inflammatory cells. Additionally, alpha-tocopherol has been shown to inhibit platelet aggregation and to enhance vasodilation.<sup>16 17</sup>

The practical application of these cardiovascular benefits were demonstrated in two studies that showed that men and women supplementing with at least 100 IU of alpha-tocopherol daily had a 37–41% drop in the risk of suboptimal heart function.<sup>18 19</sup>

Perhaps most significant is the now famous Cambridge Heart Antioxidant Study, where there was a 77% drop in suboptimal heart function in subjects who were given 400–800 IU alpha-tocopherol daily.<sup>20</sup>

Other research has shown that alpha-tocopherol may slow the progression of suboptimal memory functioning<sup>21</sup>, and may lower the risk of developing it.<sup>22</sup> In addition, research indicates that alpha-tocopherol has been successfully used for a variety of other health-related situations, including acne, seasonal pollen reactions, mutated cell growth, constipation, respiratory issues, bowel sensitivity, breast health, visual functioning, menopause, joint health, and premenstrual symptoms.<sup>23</sup>

### Gamma-tocopherol

Interestingly, the most common form of vitamin E in plants and in the American diet is gamma-tocopherol.<sup>24</sup> Although the exact role of gamma-tocopherol in human health is unclear, metabolic research suggests that it is needed for use by the body.<sup>25</sup> Some in-vitro (test tube) and animal research indicates that gamma-tocopherol may play an antioxidant role in protecting the body from free radical damage.<sup>26 27</sup> In fact, some research has indicated that gamma-tocopherol specifically protects against peroxynitrate, a powerful and harmful mutagenic radical.<sup>28</sup> Peroxynitrate is involved in creating inflammation that may further contribute to and/or be a causative factor in various disease states. This study showed that gamma-tocopherol demonstrated an ability to trap this harmful mutagen. Furthermore, gamma-tocopherol has also been shown to possess anti-inflammatory activity by blocking cyclooxygenase<sup>29</sup>, an inflammatory chemical produced in the body. Consequently, it makes sense that some research has shown that increased blood levels of gamma-tocopherol were associated with a significantly reduced risk of developing prostate cell mutations and poor cardiovascular function.<sup>30 31</sup> Some of this same research also showed similar protective effects were only seen with alpha-tocopherol and selenium when gamma-tocopherol levels were also high. Other research has shown that gamma-tocopherol assists in the excretion of sodium from the body.<sup>32</sup>

### Delta-tocopherol

Although delta-tocopherol appears to have the least vitamin E activity, it seems to provide other advantages. In-vitro research showed that delta-tocopherol was highly effective at inhibiting the growth of mutated liver cells. In comparison, gamma-tocopherol showed weak effect of inhibition and alpha-tocopherol did not show any inhibition effect at all.<sup>33</sup> In similar research on human breast cells with these three tocopherols, only delta-tocopherol was effective at inhibiting mutated cell growth.<sup>34</sup> In other research, delta-tocopherol was

shown to have the highest antioxidant activity against free radical damage in human blood, followed by gamma-tocopherol and alpha-tocopherol, respectively.<sup>35</sup>

### **Alpha-, beta- and delta-tocopherol together**

Research has shown a combination of alpha-, beta- and delta tocopherol together is more effective than alpha-tocopherol alone. For example, in one study, a mixture of these tocopherols was shown to have a stronger effect at inhibiting free radical damage to human blood cells than alpha-tocopherol alone.<sup>36</sup> The authors of this study also noted that this tocopherol mixture has been found to counteract the development of poor arterial function, whereas intake of large amounts of pure alpha-tocopherol has shown only a slight or no effect in clinical studies. In another study, the combination of tocopherols was shown to have a synergistic effect in inhibiting platelet aggregation above and beyond the effect of the individual tocopherols.<sup>37</sup>

### **Tocotrienols**

Although tocotrienols have little actual vitamin E activity, they do have significant antioxidant effects. As a matter of fact, they have better distribution in the fatty layers of the cell membranes, which makes them superior antioxidants.<sup>38 39</sup> Tocotrienols have been shown to elicit powerful antioxidant and anti-mutagenic properties, as well as supporting healthy cholesterol levels within a normal range.

In addition to their antioxidant function, tocotrienols help maintain a healthy cardiovascular system<sup>40</sup>, including protection against oxidative damage to LDL cholesterol.<sup>41</sup> In one study, many patients were able to promote healthy arterial function due to tocotrienol supplementation.<sup>42</sup> Tocotrienols have also been shown to help support healthy cholesterol levels within a normal range.<sup>43 44 45</sup>

Although a good deal of the research on tocotrienols has centered on gamma-tocotrienol, the other tocotrienols have also been shown to have valuable activity. For example, in-vitro studies indicate that the various tocotrienols may have antimutagenic benefits, particularly against breast and skin cells.<sup>46 47 48 49 50</sup> In one study, alpha-, gamma-, and delta-tocotrienols were all shown to be effective at inhibiting the growth of mutated mouse breast cells.<sup>51</sup> Another study demonstrated that alpha-, gamma-, and delta-tocotrienols helped reduce the growth of the mutated cells.<sup>52</sup> Similar results were achieved in a study where gamma- and delta-tocotrienol inhibited the growth of mutated human breast cell.<sup>53</sup>

### **Combining tocopherols and tocotrienols**

We know from the previous data presented that combining tocopherols has value over the isolated use of just alpha-tocopherol. But what happens when the tocopherols *and* the tocotrienols are combined?

In one study, a combination of tocopherols and tocotrienols were shown to significantly promote healthy cholesterol levels within a normal range in mice fed high-fat/high-cholesterol diets, compared to mice given only the combination of tocopherols with the same diet.<sup>54</sup> Clearly, the full spectrum of vitamin E isomers has greater value for promoting cardiovascular health.

In addition, research has shown that alpha-tocopherol, alpha-tocotrienol and gamma-tocopherol significantly decreased the suboptimal blood flow to the brain, while gamma-tocotrienol, delta-tocopherol and delta-tocotrienol showed no effect.<sup>55</sup> Only a combination of the isomers provides the necessary specific tocopherols and tocotrienols to provide this benefit.

### **PANTOTHENIC ACID**

Pantothenic acid is a component of coenzyme A (CoA), an essential coenzyme in a variety of reactions that sustain life. CoA is required for chemical reactions that generate energy from food (fat, carbohydrates, and proteins). The synthesis of essential fats, cholesterol, and steroid hormones requires CoA, as does the synthesis of the neurotransmitter, acetylcholine, and the hormone, melatonin. Heme, a component of hemoglobin, requires a CoA-containing compound for its synthesis. Metabolism of a number of drugs and toxins by the liver requires CoA.<sup>56</sup>

In addition, pantothenic acid is intimately involved in adrenal function, and the production of adrenal hormones associated with stress.<sup>57</sup> Experimentally-induced pantothenic acid deficiency has been associated with somnolence, fatigue, headache, paresthesia of the hands and feet followed by hyperreflexia and muscle weakness in the legs, cardiovascular instability, gastrointestinal (GI) complaints, changes in disposition, and increased susceptibility to infections.<sup>58</sup>

### **SELENIUM**

Selenium functions as a constituent of the antioxidant enzyme glutathione peroxidase, which detoxifies products of oxidized fats, and is found in the red blood cells. Selenium plays a fundamental role in regulating thyroid and other functions of the human body including reproduction, autoimmunity, glucose metabolism and bone metabolism.<sup>59</sup>

### **Selenium and immunity**

Selenium deficiency has been associated with impaired function of the immune system.<sup>60</sup> Moreover, selenium supplementation in individuals who are not overtly selenium deficient appears to stimulate the immune response. In two small studies, healthy individuals<sup>61 62</sup> and individuals with suboptimal immune response<sup>63</sup> supplemented with selenium for eight weeks showed an enhanced immune cell

response to foreign antigens compared with those taking a placebo. A considerable amount of basic research also indicates that selenium plays a role in regulating the expression of cell-signaling molecules called cytokines, which orchestrate the immune response.<sup>64</sup>

### **Selenium and mutated cell growth**

Geographic studies have consistently observed higher levels of mutated cell growth rates in populations living in areas with low soil selenium and relatively low dietary selenium intakes. Results of epidemiological studies of mutated cell growth incidence in groups with less variable selenium intakes also show a trend for individuals with lower selenium levels (blood and nails) to have a higher incidence of several different types of mutated cell growth.<sup>65</sup>

Some studies have reported that low dietary selenium intakes are associated with increased risk of mutated prostate cell growth. A case-control study within a prospective study of over 50,000 male health professionals in the U.S. found a significant inverse relationship between selenium levels and the risk mutated prostate cell growth.<sup>66</sup>

### **L-GLUTATHIONE**

Reduced glutathione (GSH) is a tripeptide consisting of L-glutamine, L-cysteine, and glycine.<sup>67</sup> It is naturally present in fruits, vegetables, and meats, and also synthesized in the body—primarily in the liver.<sup>68</sup> GSH is involved in DNA synthesis and repair, protein and prostaglandin synthesis, amino acid transport, metabolism of toxins and carcinogens, immune system function, prevention of oxidative cell damage, and enzyme activation. It is a powerful antioxidant providing protection against oxidative damage from free radicals.<sup>70 71</sup>

### **Glutathione deficiency**

Glutathione deficiency is associated with aging, age-related macular degeneration (AMD), diabetes, lung and gastrointestinal disease, pre-eclampsia, Parkinson's disease and other neurodegenerative disorders, and poor prognosis in AIDS.<sup>72 73 74 75 76 77</sup>

Oxidative stressors that can deplete GSH include ultraviolet and other radiation, viral infections, environmental toxins, household chemicals, heavy metals, surgery, inflammation, burns, septic shock, and dietary deficiencies of GSH precursors and enzyme cofactors.<sup>87 88 89 90 91 92</sup>

Replenishment of GSH is an important goal to prevent depletion and deficiency. In addition, there are some specific situations where GSH supplementation may be indicated. These situations are outlined below.

### **Detoxification**

Water soluble toxins can pass through our bodies unchanged and be eliminated in the stool, sweat or urine. Fat soluble toxins, however, cannot be excreted without undergoing metabolic transformation (detoxification) in the liver so that they can become water soluble. Liver cells have sophisticated mechanisms to break down toxic substances. These include both endogenous (produced by the body) and exogenous (obtained from the environment; i.e., xenobiotics) substances. Every drug, chemical, pesticide and hormone, is broken down or metabolized via detoxification pathways in the liver.<sup>93 94 95</sup> Phase 2 of the detoxification process involves the coupling (attaching) or conjugation of a water soluble substance which is endogenously produced or sourced by the body, to the toxin. This makes the toxic molecule more water soluble and therefore less toxic. If the molecule is large, it is then excreted via the bile. Otherwise, it is excreted in the urine.<sup>96 97</sup> GSH is one of the primary conjugating agents in phase 2, and a number of toxic molecules pass through the glutathione conjugation pathway. A deficiency of this GSH can reduce the clearance of toxins from the body.

### **Proliferation of mutated cells**

Preliminary evidence suggests GSH intake from fruits and vegetables might be associated with a reduced risk of mutated pharyngeal cell growth.<sup>98</sup> In a clinical study, patients with a proliferation of mutated colon cells were given 800 mg of GSH twice per day for at least three months. After an average of 21 weeks, 36 percent of the patients “recovered with normal diet [and] increased weight.”<sup>99</sup>

### **Seasonal infective agents**

GSH may inhibit the activity of enzymes that help the certain seasonal infective agents colonize cells lining the mouth and throat. In animal research there was lower tissue levels of the infective agents in mice fed glutathione-enriched drinking water compared to untreated mice.<sup>100</sup>

### **Chemotherapy toxicity**

Administering glutathione by intravenous injection seems to help prevent chemotherapy toxicity, improve quality of life, and provide a neuroprotective effect.<sup>101 102 103 104 105 106 107 108 109 110 111</sup>

### **Male fertility**

In a study of male fertility, glutathione was supplied in a dose of 600 mg, and administered intramuscularly for two months. In comparison to a placebo, the treatment group showed a statistically significant effect on sperm motility, specifically in the percentage of sperm demonstrating forward motility.<sup>112</sup>

### **N-ACETYL CYSTEINE**

N-acetyl cysteine (NAC) is the precursor to glutathione.<sup>113</sup> It stimulates glutathione synthesis and promotes liver detoxification; as well as acting as a powerful scavenger of free radicals.<sup>114 115</sup> Reactive oxygen species (ROS) such as hydrogen peroxide and hydroxyl-free radicals reduce intracellular and extracellular concentrations of glutathione. N-acetyl cysteine is a very efficient way to replenish glutathione and reduce damage caused by ROS.<sup>116 117</sup>

### **Effective against toxicity**

Historically the most prevalent and well-accepted use of NAC has been as an antidote for acetaminophen (Tylenol®, paracetamol) poisoning.<sup>118</sup> The resultant liver toxicity is due to an acetaminophen metabolite that depletes the liver cells of glutathione and causes liver cell damage and possibly even death. NAC has also been effective for heavy metal poisoning by gold, silver, copper, mercury, lead, and arsenic, as well as in cases of poisoning by carbon tetrachloride, acrylonitriles, halothane, paraquat, acetaldehyde, coumarin, and interferon.<sup>119</sup> Since detoxification of mercury depletes glutathione, the use of NAC doubly makes sense.

### **Beneficial for mitochondrial dysfunction**

A major cause of mitochondrial dysfunction is due to changes that take place in the respiratory chain where oxidative phosphorylation occurs. Researchers studied the effects of NAC on key elements of the respiratory chain in older rats.<sup>120</sup> After 20 weeks of treatment they found that the activities of respiratory chain were significantly higher in the treated rats compared to the controls. NAC also helped to maintain levels of the important mitochondrial antioxidant, glutathione, as well as prevented cell death in *in-vitro* studies.<sup>121</sup> In other *in-vitro* studies, NAC protected cells from programmed cell death (apoptosis) by promoting oxidative phosphorylation, mitochondrial membrane integrity, and mitochondrial homeostasis.<sup>122</sup>

### **COENZYME Q<sub>10</sub>**

Although structurally related to vitamin K, coenzyme Q<sub>10</sub> (CoQ<sub>10</sub>) is not a vitamin, but rather a coenzyme that helps to utilize oxygen as part of its important role in cellular energy metabolism. Research has also shown that CoQ<sub>10</sub> functions in a number of other beneficial ways including free radical scavenging.<sup>123</sup> Following is a brief review of that research.

### **Free radicals**

As previously described, free radicals are atoms with unpaired electrons. In the process of trying to balance itself by gaining or losing an electron, the free radical causes oxidative damage on a cellular level. CoQ<sub>10</sub> has the unique property of being able to accept or donate an electron without itself becoming a free radical. By doing this CoQ<sub>10</sub> can help neutralize free radicals and the oxidative damage they cause. This is significant since numerous

disease states are thought to be due to excessive oxidative stress of free radicals, including hydroxyl radical, peroxynitrite, superoxide anion and hydrogen peroxide. In addition, CoQ<sub>10</sub> may inhibit certain enzymes involved in the formation of these free radicals.

### **Cardiovascular support**

CoQ<sub>10</sub> has been reviewed in the scientific literature and found to be used in oral form to support cardiovascular function.<sup>124</sup>

### **Healthy heart function**

In one study, patients with suboptimal heart function experienced a significant reduction in pain, and abnormal heartbeat when supplemented with 120 mg of CoQ<sub>10</sub> daily.<sup>125</sup> In another one study, patients with suboptimal heart function were found to experience a faster loss of CoQ<sub>10</sub> during exercise.<sup>126</sup> To compensate, 150 mg of CoQ<sub>10</sub> given to patients with suboptimal heart function which not only increased their blood levels of CoQ<sub>10</sub>, but also increased their ability to exercise longer.<sup>127</sup>

A meta-analysis of eight controlled clinical trials of CoQ<sub>10</sub> supplementation in people with suboptimal heart function revealed a significant improvement in several important cardiac parameters.<sup>128</sup> Other research on CHF patients using CoQ<sub>10</sub> has shown similar benefits, including the improvement of quality of life, as well as survival.<sup>129 130</sup>

### **High blood pressure**

Research indicates that CoQ<sub>10</sub> affects blood vessels in a way that should cause support healthy blood pressure.<sup>131</sup> In fact, this has been substantiated in a number of studies where CoQ<sub>10</sub> significantly promoted healthy blood pressure within a normal range.<sup>132 133 134 135</sup> All of these studies used at least 50 mg of CoQ<sub>10</sub> taken twice daily. You should expect about 10 weeks of supplementation to pass before looking for results.

### **Mutated breast cell growth**

Biochemical, biomedical and clinical research on CoQ<sub>10</sub> and its relationship to reducing the growth of mutated breast cells has evolved internationally over 35 years. In published research, patients were able to reduce the spread of mutated breast cell growth by supplementing with 390 mg of CoQ<sub>10</sub> daily.<sup>136</sup> This was also seen when CoQ<sub>10</sub> was used at a lower doses of 90 mg daily in combination with other antioxidants including vitamin C, Vitamin E, beta-carotene, selenium, and essential fatty.<sup>137</sup> Without the benefit of other antioxidants, however the 390 mg dose of CoQ<sub>10</sub>.<sup>138</sup>

### **Blood glucose levels**

Research has shown that some individuals with higher blood glucose levels may have a deficiency of CoQ<sub>10</sub>, which may be further exacerbated by the use

of certain commonly used drugs. Such a deficiency of CoQ10 in the pancreas could impair aspects of energy metabolism and the biosynthesis of insulin.<sup>139</sup>

### **Periodontal disease**

Did you know that in Japan over half the dentists recommend supplements of CoQ10 to promote healthy gums? There are very good reasons for this. First of all, research has shown that patients with suboptimal gum health have a CoQ10 deficiency of in their gums.<sup>140</sup> Secondly, treatment with CoQ10 in these significant improvement in the condition of the gums.<sup>141</sup> The reason that CoQ10 works may have to do with an ability to inhibit bacterial growth due to improved oxygen metabolism at the cellular level (bacteria often do not survive in the presence of oxygen).

### **ALPHA LIPOIC ACID**

Alpha Lipoic Acid (ALA), also known as thioctic acid, has gained considerable attention as an antioxidant. ALA combats particularly nasty free radicals such as superoxide radicals, hydroxyl radicals, hypochlorous acid, peroxy radicals, and singlet oxygen, thereby reducing oxidative stress. ALA is a small molecule, soluble in both water and fat. This allows it to work both inside the cell and at the membrane level, making ALA a particularly valuable antioxidant.

### **Oxidative stress and ALA**

In general, antioxidants provide a preventive measure against the hazards of oxidative stress with their ability to neutralize, balance and sponge up free radicals by coupling with unpaired electrons. ALA is unique since it not only acts as antioxidant against free radicals, but it also helps in the regeneration process of expended antioxidants, returning them to their reduced, antioxidant-potent form. In other words, it helps to recycle other antioxidants. Consequently, ALA is a key ingredient in providing the maximum interaction among antioxidants, enhancing their cell protection abilities. It has the unique ability to enhance the power of vitamins E and C and Glutathione. This results in an antioxidant network which provides more complete cell protection from destructive free radicals.<sup>142</sup> In animal research, ALA reversed a number of age-related changes in the brains of rats, including increased activity of antioxidant enzymes and decreased production of free radicals.<sup>143</sup>

### **The metabolic antioxidant**

In addition to its antioxidant functions, ALA also has a metabolic role to play in the body: It facilitates the production of energy by aiding in the metabolism of glucose, and in the mitochondrial Krebs cycle which produces adenosine triphosphate (ATP, the energy currency of the body).<sup>144</sup> This energy production is vital for athletes and active individuals to function at maximum efficiency, or peak levels. Since ALA has

both antioxidant and metabolic functions, it is considered to be a metabolic antioxidant.

### **Other uses for ALA**

In his book, *The Wrinkle Cure*, Dr. Nicholas Perricone<sup>145</sup> recommends ALA as part of a strategy for maintaining healthy skin and reducing the potential for wrinkles. ALA also has value as part of a complete program for facilitating the liver's detoxification process.<sup>146</sup> In research, 150 mg of ALA per day for one month was shown to improve visual function in people with inner ocular pressure.<sup>147</sup> Finally, in a preliminary trial<sup>148</sup> three patients with suboptimal liver function and dilated veins in the esophagus received a combination of 300 mg ALA twice daily, 300 mg silimarin from milk thistle three times daily, and 200 mg selenium twice daily). After five to eight months, all three patients had significant improvements in their liver function and overall health (note: they also used other supportive supplements which included vitamin C and B vitamins).

### **REFERENCES**

- <sup>1</sup> Omenn GS. Chemoprevention of lung cancer: the rise and demise of beta-carotene. *Annu Rev Public Health* 1998;19:73-99.
- <sup>2</sup> Erlinger TP, Guallar E, Miller ER, et al. Relationship between systemic markers of inflammation and serum beta-carotene levels. *Arch Intern Med* 2001;161:1903-8.
- <sup>3</sup> van der Vliet A. Cigarettes, cancer and the carotenoids: a continuing, unresolved antioxidant paradox. *Am J Clin Nutr* 2000;72:1421-3.
- <sup>4</sup> Green A, Williams G, Neale R, et al. Daily sunscreen applications and beta-carotene supplementation in prevention of basal-cell and squamous-cell carcinomas of the skin: a randomized controlled trial. *Lancet* 1999;354:723-9.
- <sup>5</sup> Shekelle RB, Liu S, Raynor WJ Jr, Rossof AH, Paul O, Shryock AM, Stamler J. Dietary vitamin A and risk of cancer in the Western Electric Study. *Lancet* 1981; 2:1185-90.
- <sup>6</sup> Stich HF, Rosin MP, Hornby AP, et al. Remission of oral leukoplakias and micronuclei in tobacco/betel quid chewers treated with beta-carotene and with beta-carotene plus vitamin A. *Int J Cancer* 1988;42:195-9
- <sup>7</sup> Garewal HS, Katz RV, Meyskens F, Pitcock J, Morse D, Friedman S, Peng Y, Pendrys DG, Mayne S, Alberts D, Kiersch T, Graver E. β-Carotene produces sustained remission in patients with oral leukoplakia. *Arch Otolaryngol Head Neck Surg* 1999;125:1305-10.
- <sup>8</sup> McKevooy GK, ed. AHFS Drug Information. Bethesda, MD: American Society of Health-System Pharmacists, 1998.
- <sup>9</sup> Leibovitz B, Siegel BV. Ascorbic acid and the immune response. *Adv Exp Med Biol* 1981;135:1-25.
- <sup>10</sup> Levine M, Rumsey SC, Daruwala R, et al. Criteria and recommendations for vitamin C intake. *JAMA* 1999;281:1415-23.
- <sup>11</sup> Kaufmann PA, Gnecci-Ruscione T, di Terlizzi M, et al. Coronary heart disease in smokers: vitamin C restores coronary microcirculatory function. *Circulation* 2000;102:1233-8.
- <sup>12</sup> Traber MG. Utilization of vitamin E. *Biofactors*. 1999;10(2-3):115-120.
- <sup>13</sup> Seager SL, Slabaugh MR. *Chemistry for Today: General, Organic, and Biochemistry*, Fourth Edition. Pacific Grove, California: Brooks/Cole; 2000:364.
- <sup>14</sup> Traber MG. Vitamin E. In: Shils ME, Shike M, Ross AC, Caballero B, Cousins RJ, eds. *Modern Nutrition in Health and Disease*. Philadelphia: Lippincott Williams & Wilkins; 2006:396-411.
- <sup>15</sup> Bruno RS, Leonard SW, Atkinson J, et al. Faster plasma vitamin E disappearance in smokers is normalized by vitamin C supplementation. *Free Radic Biol Med*. 2006;40(4):689-697.

- <sup>16</sup> Food and Nutrition Board, Institute of Medicine. Vitamin E. Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. Washington D.C.: National Academy Press; 2000:186-283.
- <sup>17</sup> Traber MG. Does vitamin E decrease heart attack risk? summary and implications with respect to dietary recommendations. *J Nutr*. 2001;131(2):395S-397S.
- <sup>18</sup> Rimm EB, Stampfer MJ, Ascherio A, et al. Vitamin E consumption and the risk of coronary heart disease in men. *N Engl J Med* 1993; 328:1450-6.
- <sup>19</sup> Stampfer MJ, Hennekens CH, Manson JE, et al. Vitamin E consumption and the risk of coronary heart disease in women. *N Engl J Med* 1993; 328:1444-9.
- <sup>20</sup> Stephens NG, Parsons A, Schofield PM, et al. Randomised controlled trial of vitamin E in patients with coronary disease: Cambridge Heart Antioxidant Study (CHAOS). *Lancet* 1996; 347:781-6.
- <sup>21</sup> Sano M, Ernesto C, Thomas RG, et al. A controlled trial of selegiline, alpha-tocopherol, or both as treatment for Alzheimer's disease. *N Engl J Med* 1997; 336:1216-22.
- <sup>22</sup> Morris MC, Beckett LA, Scherr PA, et al. Vitamin E and vitamin C supplement use and risk of incident Alzheimer disease. *Alzheimer Dis Assoc Disord* 1998;12:121-6.
- <sup>23</sup> Bruno G. *Ailments & Natural Remedies*. New York: Fifty-Ninth Street Bridge Publishing Corp; 2001.
- <sup>24</sup> Jiang Q, Christen S, Shigenaga MK, Ames BN.  $\gamma$ -Tocopherol, the major form of vitamin E in the US diet, deserves more attention. *Am J Clin Nutr* 2001;74:714-22.
- <sup>25</sup> Traber MG, Elsner A, Brigelius-Flohe R. Synthetic as compared with natural vitamin E is preferentially excreted as alpha-CEHC in human urine: studies using deuterated alpha-tocopheryl acetates. *FEBS Lett* 1998;437(1-2):145-148.
- <sup>26</sup> Christen S, Woodall AA, Shigenaga MK, Southwell-Keely PT, Duncan MW, Ames BN. gamma-tocopherol traps mutagenic electrophiles such as NO(X) and complements alpha-tocopherol: physiological implications. *Proc Natl Acad Sci U S A*. 1997;94(7):3217-3222.
- <sup>27</sup> Li D, Saldeen T, Mehta JL. gamma-tocopherol decreases ox-LDL-mediated activation of nuclear factor-kappaB and apoptosis in human coronary artery endothelial cells. *Biochem Biophys Res Commun*. 1999;259(1):157-161.
- <sup>28</sup> Christen S, Woodall AA, Shigenaga MK, et al. g-Tocopherol traps mutagenic electrophiles such as NOx and complements a-tocopherol: Physiological implications (antimutagenylipid peroxidation/low-density lipoprotein/inflammation). *Proc Natl Acad Sci* 1997; 94:3217-3222.
- <sup>29</sup> Jiang Q, Christen S, Shigenaga MK, Ames BN.  $\gamma$ -Tocopherol, the major form of vitamin E in the US diet, deserves more attention. *Am J Clin Nutr* 2001;74:714-22.
- <sup>30</sup> Helzlsouer KJ, Huang HY, Alberg AJ, et al. Association between alpha-tocopherol, gamma-tocopherol, selenium, and subsequent prostate cancer. *J Natl Cancer Inst*. 2000;92(24):2018-2023.
- <sup>31</sup> Jiang Q, Christen S, Shigenaga MK, Ames BN.  $\gamma$ -Tocopherol, the major form of vitamin E in the US diet, deserves more attention. *Am J Clin Nutr* 2001;74:714-22.
- <sup>32</sup> Brigelius-Flohe R, Traber MG. Vitamin E: function and metabolism. *FASEB J* 1999; 13:1145-1155.
- <sup>33</sup> Junxia M, Junsheng G, Faji Z, Donglian C. [Effect of apoptosis induced by different vitamin E homologous analogues in human hepatoma cells (HepG2)] Wei sheng yan jiu = Journal of hygiene research 2003; 32(2):131-3.
- <sup>34</sup> Yu W, Simmons-Menchaca M, Gapor A, et al. Induction of apoptosis in human breast cancer cells by tocopherols and tocotrienols. *Nutrition and cancer* 1999; 33(1):26-32.
- <sup>35</sup> Pierre S, Jean-Francois L, Nicolas V, et al. ESR study of a biological assay on whole blood: antioxidant efficiency of various vitamins. *Biochimica et biophysica acta* 2003; 1621(1):1-8.
- <sup>36</sup> Meilin L; Rolf W, Agneta W, Tom S. Mixed tocopherols have a stronger inhibitory effect on lipid peroxidation than alpha-tocopherol alone. *Journal of cardiovascular pharmacology* 2002; 39(5):714-21.
- <sup>37</sup> Li D, Saldeen T, Romeo F, Mehta JL. Different isoforms of tocopherols enhance nitric oxide synthase phosphorylation and inhibit human platelet aggregation and lipid peroxidation: implications in therapy with vitamin E. *Journal of cardiovascular pharmacology and therapeutics* 2001; 6(2):155-61.
- <sup>38</sup> Kamal-Eldin A, Appelqvist LA. The chemistry and antioxidant properties of tocopherols and tocotrienols. *Lipids* 1996; 31:671-701.
- <sup>39</sup> Kamat JP, Devasagayam TPA. Tocotrienols from palm oil as potent inhibitors of lipid peroxidation and protein oxidation in rat brain mitochondria. *Neuroscience Letters* 1995; 195:179-82.
- <sup>40</sup> Theriault A, Chao JT, Wang Q, et al. A review of its therapeutic potential. *Clin Biochem* 1999;32:309-19.
- <sup>41</sup> Suarna C, Hood RL, Dean RT, Stocker R. Comparative antioxidant activity of tocotrienols and other natural lipid-soluble antioxidants in a homogeneous system, and in rat and human lipoproteins. *Biochem Biophys Acta* 1993; 1166:163-70.
- <sup>42</sup> Tomeo AC, Geller M, Watkins TR, et al. Antioxidant effects of tocotrienols in patients with hyperlipidemia and carotid stenosis. *Lipids* 1995; 30:1179-83.
- <sup>43</sup> Parker RA, Pearce, BC, et al. Tocotrienols Decrease Cholesterol Synthesis in HEPG2 Cells by A Novel Post-Transcriptional Suppression of HMG-CoA Reductase. *FASEB J* 1990;4:1744A.
- <sup>44</sup> Qureshi AA, Bradlow BA, Brace L, et al. Response of hypercholesterolemic subjects to administration of tocotrienols. *Lipids* 1995; 30:1171-7.
- <sup>45</sup> Qureshi AA, Qureshi N, Wright JJ, et al. Lowering of serum cholesterol in hypercholesterolemic humans by tocotrienols (palmvitae). *Am J Clin Nutr* 1991; 53:1021S-1026S.
- <sup>46</sup> Goh SH, Hew NF, Norhanom AW, Yadav M. Inhibition of tumour promotion by various palm-oil tocotrienols. *Int J Cancer* 1994; 57:529-31.
- <sup>47</sup> Yu W, Simmons-Menchaca M, Gapor A, et al. Induction of apoptosis in human breast cancer cells by tocopherols and tocotrienols. *Nutr Cancer* 1999; 33:26-32.
- <sup>48</sup> Mo H, Elson CE. Apoptosis and cell-cycle arrest in human and murine tumor cells are initiated by isoprenoids. *J Nutr* 1999; 129:804-13.
- <sup>49</sup> Nesaretnam K, Stephen R, Dils R, Darbre P. Tocotrienols inhibit the growth of human breast cancer cells irrespective of estrogen receptor status. *Lipids* 1998; 33:461-9.
- <sup>50</sup> He L, Mo H, Hadisusilo S, et al. Isoprenoids suppress the growth of murine B16 melanomas in vitro and in vivo. *J Nutr* 1997; 127:668-74.
- <sup>51</sup> McIntyre BS, Briski KP, Tirmenstein MA, et al. Antiproliferative and apoptotic effects of tocopherols and tocotrienols on normal mouse mammary epithelial cells. *Lipids* 2000; 35(2):171-80.
- <sup>52</sup> Guthrie N, Gapor A, Chambers AF, Carroll KK. Inhibition of proliferation of estrogen receptor-negative MDA-MB-435 and -positive MCF-7 human breast cancer cells by palm oil tocotrienols and tamoxifen, alone and in combination. *Journal of nutrition* 1997; 127(3):544S-548S.
- <sup>53</sup> Nesaretnam K, Stephen R, Dils R, Darbre P. Tocotrienols inhibit the growth of human breast cancer cells irrespective of estrogen receptor status. *Lipids* 1998; 33:461-9.
- <sup>54</sup> Black TM, Wang P, Maeda N, Coleman RA. Palm tocotrienols protect ApoE +/- mice from diet-induced atheroma formation. *Journal of nutrition* 2000; 130(10):2420-6.
- <sup>55</sup> Kenichi M, Takamitsu T, Fengling P, et al. Vitamin E isoforms alpha-tocotrienol and gamma-tocopherol prevent cerebral infarction in mice. *Neurosci Lett* 2003 30;337(1):56-60.
- <sup>56</sup> Brody T. *Nutritional Biochemistry*. 2nd ed. San Diego: Academic Press; 1999.
- <sup>57</sup> Kutsky R, *Handbook of Vitamins and Hormones*. New York: Van Nostrand Reinhold Company; 1973:208.
- <sup>58</sup> McKeovoy GK, ed. *AHFS Drug Information*. Bethesda, MD: American Society of Health-System Pharmacists, 1998.
- <sup>59</sup> Kaprara A, Krassas GE. Selenium and thyroidal function; the role of immunoassays. *Hell J Nucl Med* 2006;9(3):195-203.
- <sup>60</sup> McKenzie RC, Beckett GJ, Arthur JR. Effects of selenium on immunity and aging. In: Hatfield DL, Berry MJ, Gladyshev VN, eds. *Selenium: Its Molecular Biology and Role in Human Health*. 2nd ed. New York: Springer; 2006:311-323.
- <sup>61</sup> Roy M, Kiremidjian-Schumacher L, Wishe HI, Cohen MW, Stotzky G. Supplementation with selenium and human immune cell functions. I. Effect on lymphocyte proliferation and interleukin

- 2 receptor expression. *Biol Trace Elem Res.* 1994;41(1-2):103-114.
- <sup>62</sup> Kiremidjian-Schumacher L, Roy M, Wishe HI, Cohen MW, Stotzky G. Supplementation with selenium and human immune cell functions. II. Effect on cytotoxic lymphocytes and natural killer cells. *Biol Trace Elem Res.* 1994;41(1-2):115-127.
- <sup>63</sup> Kiremidjian-Schumacher L, Roy M, Glickman R, et al. Selenium and immunocompetence in patients with head and neck cancer. *Biol Trace Elem Res.* 2000;73(2):97-111.
- <sup>64</sup> Baum MK, Miguez-Burbano MJ, Campa A, Shor-Posner G. Selenium and interleukins in persons infected with human immunodeficiency virus type 1. *J Infect Dis.* 2000;182 Suppl 1:S69-73.
- <sup>65</sup> Garland M, Morris JS, Stampfer MJ, et al. Prospective study of toenail selenium levels and cancer among women. *J Natl Cancer Inst.* 1995;87(7):497-505.
- <sup>66</sup> Yoshizawa K, Willett WC, Morris SJ, et al. Study of prediagnostic selenium level in toenails and the risk of advanced prostate cancer. *J Natl Cancer Inst.* 1998;90(16):1219-1224.
- <sup>67</sup> Glutathione, reduced (GSH). Monograph. *Altern Med Rev* 2001;6(6):601-7.
- <sup>68</sup> Anderson ME. Glutathione: an overview of biosynthesis and modulation. *Chem Biol Interact* 1998;24;111-112:1-14.
- <sup>69</sup> Lu SC. Regulation of hepatic glutathione synthesis: current concepts and controversies. *FASEB J* 1999;13:1169-83.
- <sup>70</sup> Lomaestro BM, Malone M. Glutathione in health and disease: pharmacotherapeutic issues. *Ann Pharmacother* 1995;29:1263-73.
- <sup>71</sup> Powers SK, Hamilton K. Antioxidants and exercise. *Clin Sports Med* 1999;18:525-36.
- <sup>72</sup> Lomaestro BM, Malone M. Glutathione in health and disease: pharmacotherapeutic issues. *Ann Pharmacother* 1995;29:1263-73.
- <sup>73</sup> Walsh SW, Wang Y. Deficient glutathione peroxidase activity in preeclampsia is associated with increased placental production of thromboxane and lipid peroxides. *Am J Obstet Gynecol* 1993;169:1456-61.
- <sup>74</sup> Knapen MF, Mulder TP, Van Rooij IA, et al. Low whole blood glutathione levels in pregnancies complicated by preeclampsia or the hemolysis, elevated liver enzymes, low platelets syndrome. *Obstet Gynecol* 1998;92:1012-5.
- <sup>75</sup> Knapen MF, Peters WH, Mulder TP, et al. Glutathione and glutathione-related enzymes in decidua and placenta of controls and women with pre-eclampsia. *Placenta* 1999;20:541-6.
- <sup>76</sup> Marshall KA, Reist M, Jenner P, et al. The neuronal toxicity of sulfite plus peroxynitrite is enhanced by glutathione depletion: implications for Parkinson's disease. *Free Radic Biol Med* 1999;27:515-20.
- <sup>77</sup> Jenner P, Olanow CW. Understanding cell death in Parkinson's disease. *Ann Neurol* 1998;44(3 Suppl 1):S72-84.
- <sup>78</sup> Jenner P. Oxidative mechanisms in nigral cell death in Parkinson's disease. *Mov Disord* 1998;(13 Suppl 1):24-34.
- <sup>79</sup> Pearce RK, Owen A, Daniel S, et al. Alterations in the distribution of glutathione in the substantia nigra in Parkinson's disease. *J Neural Transm* 1997;104:661-77.
- <sup>80</sup> Merad-Boudia M, Nicole A, Santiard-Baron D, et al. Mitochondrial impairment as an early event in the process of apoptosis induced by glutathione depletion in neuronal cells: relevance to Parkinson's disease. *Biochem Pharmacol* 1998;56:645-55.
- <sup>81</sup> Sechi G, Deledda MG, Bua G, et al. Reduced intravenous glutathione in the treatment of early Parkinson's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 1996;20:1159-70.
- <sup>82</sup> Droge W, Holm E. Role of cysteine and glutathione in HIV infection and other diseases associated with muscle wasting and immunological dysfunction. *FASEB J* 1997;11:1077-89.
- <sup>83</sup> Herzenberg LA, De Rosa SC, Dubs JG, et al. Glutathione deficiency is associated with impaired survival in HIV disease. *Proc Natl Acad Sci USA* 1997;94:1967-72.
- <sup>84</sup> Samiec PS, Drews-Botsch C, Flagg EW, et al. Glutathione in human plasma: decline in association with aging, age-related macular degeneration, and diabetes. *Free Radic Biol Med* 1998;24:699-704.
- <sup>85</sup> Bains JS, Shaw CA. Neurodegenerative disorders in humans: the role of glutathione in oxidative stress-mediated neuronal death. *Brain Res Brain Res Rev* 1997;25:335-58.
- <sup>86</sup> Loguercio C, Di Pierro M. The role of glutathione in the gastrointestinal tract: a review. *Ital J Gastroenterol Hepatol* 1999;31:401-7.
- <sup>87</sup> Cai J, Nelson KC, Wu M, et al. Oxidative damage and protection of the RPE. *Progr Retinal Eye Res* 2000;19:205-221.
- <sup>88</sup> Kidd PM. Glutathione: systemic protectant against oxidative and free radical damage. *Altern Med Rev* 1997;1:155-176.
- <sup>89</sup> Look MP, Rockstroh JK, Rao GS, et al. Serum selenium, plasma glutathione (GSH) and erythrocyte glutathione peroxidase (GSH-Px)-levels in asymptomatic versus symptomatic human immunodeficiency virus-1 (HIV-1)-infection. *Eur J Clin Nutr* 1997;51:266-272.
- <sup>90</sup> Luo J-L, Hammarqvist F, Andersson K, et al. Surgical trauma decreases glutathione synthetic capacity in human skeletal muscle tissue. *Am J Physiol* 1998;275:E359-E365.
- <sup>91</sup> Spies CD, Reinhart K, Witt I, et al. Influence of N-acetylcysteine on direct indicators of tissue oxygenation in septic shock patients: results from a prospective, randomized, double-blind study. *Crit Care Med* 1994;22:1738-1746.
- <sup>92</sup> Whitcomb DC, Block GD. Association of acetaminophen hepatotoxicity with fasting and ethanol use. *JAMA* 1994;272:1845-1850.
- <sup>93</sup> Murray Rk, Granner DK, Mayes PA, Rodwell VW. Harper's Biochemistry, 25<sup>th</sup> ed. New York: McGraw Hill; 200:780-786.
- <sup>94</sup> Lüllmann H. Mohr K, Ziegler A, Bieger D. Color Atlas of Pharmacology, 2<sup>nd</sup> ed. Stuttgart: Thieme; 2000:32-39.
- <sup>95</sup> Roundtree R. The Use of Phytochemicals in the Biotransformation and Elimination of Environmental Toxins. IN Medicines from the Earth 2003: Official Proceedings. Brevard, North Carolina: Gaia Herbal Research Institute; 2003:115-128.
- <sup>96</sup> Murray Rk, Granner DK, Mayes PA, Rodwell VW. Harper's Biochemistry, 25<sup>th</sup> ed. New York: McGraw Hill; 200:780-786.
- <sup>97</sup> Lüllmann H. Mohr K, Ziegler A, Bieger D. Color Atlas of Pharmacology, 2<sup>nd</sup> ed. Stuttgart: Thieme; 2000:32-39.
- <sup>98</sup> Flagg EW, Coates RJ, Jones DP, et al. Dietary glutathione intake and the risk of oral and pharyngeal cancer. *Am J Epidemiol* 1994;139:453-65.
- <sup>99</sup> Garcia-Girlat E, Perdereau B, Brixy F, et al. Preliminary study of glutathione, L-cysteine and anthocyanins (Recancostat Compositum™) in metastatic colorectal carcinoma with malnutrition. Seventh International Congress on Anti-cancer Treatment, February 3–6, 1997, Palai des Congrès, Paris, France.
- <sup>100</sup> Cai J, Chen Y, Seth S, Furukawa S, Compans RW, Jones DP. Inhibition of influenza infection by glutathione. *Free Radic Biol Med* 2003;34(7):928-36.
- <sup>101</sup> Smyth JF, Bowman A, Perren T, et al. Glutathione reduces the toxicity and improves quality of life of women diagnosed with ovarian cancer treated with cisplatin: results of a double-blind, randomised trial. *Ann Oncol* 1997;8:569-73.
- <sup>102</sup> Cascinu S, Cordella L, Del Ferro E, et al. Neuroprotective effect of reduced glutathione on cisplatin-based chemotherapy in advanced gastric cancer: a randomized, double-blind, placebo-controlled trial. *J Clin Oncol* 1995;13:26-32.
- <sup>103</sup> Links M, Lewis C. Chemoprotectants: a review of their clinical pharmacology and therapeutic efficacy. *Drugs* 1999;57:293-308.
- <sup>104</sup> Leone R, Fracasso ME, Soresi E, et al. Influence of glutathione administration on the disposition of free and total platinum in patients after administration of cisplatin. *Cancer Chemother Pharmacol* 1992;29:385-90.
- <sup>105</sup> Graziano F, Cardarelli N, Marcellini M, et al. A pilot clinical trial of postoperative intensive weekly chemotherapy using cisplatin, epi-doxorubicin, 5-fluorouracil, 6S-leucovorin, glutathione and filgrastim in patients with resected gastric cancer. *Tumori* 1998;84:368-71.
- <sup>106</sup> Plaxe S, Freddo J, Kim S, et al. Phase I trial of cisplatin in combination with glutathione. *Gynecol Oncol* 1994;55:82-6.
- <sup>107</sup> Locatelli MC, D'Antona A, Labianca R, et al. A phase II study of combination chemotherapy in advanced ovarian carcinoma with cisplatin and cyclophosphamide plus reduced glutathione as potential protective agent against cisplatin toxicity. *Tumori* 1993;79:37-9.
- <sup>108</sup> Di Re F, Bohm S, Oriana S, et al. High-dose cisplatin and cyclophosphamide with glutathione in the treatment of advanced ovarian cancer. *Ann Oncol* 1993;4:55-61.

- <sup>109</sup> Parnis FX, Coleman RE, Harper PG, et al. A randomised double-blind placebo controlled clinical trial assessing the tolerability and efficacy of glutathione as an adjuvant to escalating doses of cisplatin in the treatment of advanced ovarian cancer. *Eur J Cancer* 1995;31A:1721.
- <sup>110</sup> Cascinu S, Frontini L, Comella G, et al. Intensive weekly chemotherapy is not effective in advanced pancreatic cancer patients: a report from the Italian Group for the Study of Dig. Tract Cancer (GISCAD). *Br J Cancer* 1999;79(3-4):491-4.
- <sup>111</sup> Cascinu S, Labianca R, Alessandrini P, et al. Intensive weekly chemotherapy for advanced gastric cancer using fluorouracil, cisplatin, epi-doxorubicin, 6S-leucovorin, glutathione, and filgrastim: a report from the Italian Group for the Study of Digestive Tract Cancer. *J Clin Oncol* 1997;15:3313-9.
- <sup>112</sup> Lenzi A, Culasso F, Gandini L, et al. Placebo-controlled, double blind, cross-over trial of glutathione therapy in male infertility. *Hum Reprod* 1993; 8:1657-1662.
- <sup>113</sup> Anonymous. *Alternative Medicine Review* 2000; 5(5); 467-471.
- <sup>114</sup> De Vries N, De Flora S. N-Acetyl-L-Cysteine. *J Cell Biochem* 1993;17F:S270-S277.(-)
- <sup>115</sup> De Flora S, Bennicelli C, Camoirano A, et al. In vivo effects of N-acetylcysteine on glutathione metabolism and on the biotransformation of carcinogenic and/or mutagenic compounds. *Carcinogenesis* 1985;6:1735-1745.
- <sup>116</sup> Weinbroum AA, Rudick V, Ben-Abraham R, Karchevski E. N-acetyl-L-cysteine for preventing lung reperfusion injury after liver ischemia-reperfusion: a possible dual protective mechanism in a dose-response study. *Transplantation* 2000;69:853-9.
- <sup>117</sup> Kelly GS. Clinical applications of N-acetylcysteine. *Altern Med Rev* 1998;3:114-27.
- <sup>118</sup> Anonymous. *Alternative Medicine Review* 2000; 5(5); 467-471.
- <sup>119</sup> Zimet I. Acetylcysteine: A drug that is much more than a mucokinetic. *Biomed & Pharmacother* 1988;42:513-520.
- <sup>120</sup> Chakraborti S, Batabyal S, Ghosh S, Chakraborti T. Protective role of N-acetylcysteine against the age-related decline in oxidative phosphorylation in pulmonary smooth muscle mitochondria. *Med Sci Res* 1999; 27(1):39-40.
- <sup>121</sup> Banachocha M. Therapeutic potential of N-acetylcysteine in age-related mitochondrial neurodegenerative diseases. *Medical Hypotheses* 2001; 56(4):472-477.
- <sup>122</sup> Cossarizza A, Franceschi C, Monti D, et al. Protective effect of N-Acetylcysteine in tumor necrosis factor-alpha-induced apoptosis in U937 cells: The role of mitochondria. *Experimental Cell Research* 1995; 220:232-240.
- <sup>123</sup> Pepping J, *Am J Health-Syst Pharm* (1999) 56:519-521.
- <sup>124</sup> Greenberg S, Frishman WH, *J Clin Pharmacol* (1990) 30(7):596-608.
- <sup>125</sup> Singh RB, et al, *Cardiovasc Drugs Ther* (1998) 12(4):347-53.
- <sup>126</sup> Karlsson J, et al, *Ann Med* (1991) 23(3):339-44.
- <sup>127</sup> Kamikawa T, *Am J Cardiol* (1985) 56 (4):247-51.
- <sup>128</sup> Soja AM, Mortensen SA, *Ugeskr Laeger* (1997) 159(49):7302-8.
- <sup>129</sup> Munkholm H, Hansen HH, Rasmussen K, *Biofactors* (1999) 9(2-4):285-9.
- <sup>130</sup> Sinatra ST, *Conn Med* (1997) 61(11):707-11.
- <sup>131</sup> Digiesi V, et al, *Curr Ther Res* (1992) 51:668-72.
- <sup>132</sup> Folkers K, et al, *Res Commun Chem Pathol Pharmacol* (1981) 31:129-40.
- <sup>133</sup> Langsjoen P, et al, *Mol Aspects Med* (1994) 15(Suppl):S265-72.
- <sup>134</sup> Digiesi V, et al, *Molec Aspects Med* (1994) 15(Suppl):S257-63.
- <sup>135</sup> Digiesi V, Cantini F, Brodbeck B, *Curr Ther Res* (1990) 47:841-45.
- <sup>136</sup> Lockwood K, et al, *Biochem Biophys Res Commun* (1995) 212(1):172-7.
- <sup>137</sup> Lockwood K, et al, *Mol Aspects Med* (1994) 15 Suppl:s231-40.
- <sup>138</sup> Lockwood K, Moesgaard S, Folkers K, *Biochem Biophys Res Commun* (1994) 199(3):p1504-8
- <sup>139</sup> Kishi T, et al, *J Med* (1976) 7(3-4):307-21.
- <sup>140</sup> Hansen IL, et al, *Res Commun Chem Pathol Pharmacol* (1976) 14(4):729-38.
- <sup>141</sup> Wilkinson EG, Arnold RM, Folkers K, *Res Commun Chem Pathol Pharmacol* (1976) 14(4):715-9.
- <sup>142</sup> Packer, L., et al, *Free Radic Biol Med* 1995; 19(2): 227-50.
- <sup>143</sup> Martin D, Towey M, Horrobin D, Lynch M. A diet enriched in alpha lipoic acid reverses the age-related compromise in antioxidant defenses in rat cortical tissue. *Nutr Neurosci* 2000;3(3):193-206.
- <sup>144</sup> Beitner H. Randomized, placebo controlled, double-blind study on the clinical efficacy of a cream containing 5% alpha-lipoic acid related to photoaging of facial skin. *Br J Dermatol* 2003;149:841-9.
- <sup>145</sup> Perricone, N. *The Wrinkle Cure*. Rodale; 2000.
- <sup>146</sup> Roundtree R. The Use of Phytochemicals in the Biotransformation and Elimination of Environmental Toxins. IN *Medicines from the Earth 2003: Official Proceedings*. Brevard, North Carolina: Gaia Herbal Research Institute; 2003:115-128.
- <sup>147</sup> Filina AA, Davydova NG, Endrikhovskii SN, et al. Lipoic acid as a means of metabolic therapy of open-angle glaucoma. *Vestn Oftalmol* 1995;111:6-8.
- <sup>148</sup> Berkson BM. A conservative triple antioxidant approach to the treatment of hepatitis C. Combination of Alpha lipoic acid (thioctic acid), silymarin, and selenium: three case histories. *Med Klin* 1999; 94 Suppl 3:84-9.



For more than two decades, Huntington College of Health Sciences (HCHS) has offered more than a conventional undergraduate or graduate education. Our accredited\*, distance learning degrees and diploma programs also include the breadth of responsible complementary and alternative medicine viewpoints, providing our students with a well-rounded and comprehensive approach to nutrition and the health sciences:

- Master of Science in Nutrition
- Bachelor of Health Science in Nutrition
- Associate of Science in Applied Nutrition
- Diploma in Comprehensive Nutrition
- Diploma in Dietary Supplement Science
- Diploma in Sports Nutrition
- Diploma in Women's Nutrition
- Diploma in Natural Sciences
- Diploma in Small Business Management

1204D Kenesaw  
Knoxville, TN 37919  
865-524-8079 • 800-290-4226  
E-Mail: [studentservices@hchs.edu](mailto:studentservices@hchs.edu)  
[www.hchs.edu](http://www.hchs.edu)

\*Accredited member Distance Education & Training Council.